

Manure nutrient management planning has moved into the spotlight in recent years. While crop producers have traditionally considered crop nutrients supplied by manure, increased attention to factors influencing manure nutrient use can improve overall crop production efficiency and profitability. Increasing environmental concerns for surface water and groundwater, and accelerated concentration of livestock production also have increased the need for improved manure nutrient planning. Managing manure for efficient crop production, while minimizing potential environmental concerns, is more complex than simply disposing of the manure on agricultural lands. It entails more than focusing on the total amount of nutrients contained in manure.

A significant portion of nutrients in manure are in organic forms and not available for plant uptake until undergoing mineralization. Mineralization is the conversion of organic forms of nutrients to plant-available, inorganic forms of nutrients by soil microbes during decomposition of organic materials such as manure. Because soil microbial activity is affected by uncontrollable factors such as soil temperature and moisture, crediting the amount of crop-available nutrients provided by manure application cannot accurately be predicted. Additionally, the analysis of the specific manure used, method of application, time delay between manure application and incorporation (if incorporated), calibration of application equipment (rate and uniformity), and other specific field/crop conditions all affect the amount of manure nutrients that will become available to a crop. Unless all these factors are considered, the crop nutrients in the manure will not be used most efficiently and questions pertaining to environmental stewardship may remain.

## Environmental Considerations

Manure represents a valuable source of nitrogen (N), phosphorus (P), potassium (K), and other crop nutrients that should be considered carefully when developing an overall crop nutrient management plan. With improper management, manure applications pose potential environmental risks. Manure typically contains relatively large amounts of nitrogen. Overapplication can result in high levels of residual soil nitrate-N, posing risk to groundwater supplies through leaching.

Proper crediting of estimated nitrogen availability from manure is important to minimize potential problems.


Additionally, manure is an important source of phosphorus, and proper manure application management and nutrient crediting is important in protecting water quality. Non-point phosphorus runoff from agricultural fields is associated with over-enrichment (termed “eutrophication”) of some surface water bodies. Eutrophication can result in excessive algal and aquatic weed growth that depletes oxygen in these waters, resulting in fish kills and the need for affected waters to be treated by public water systems. Eutrophication also has harmful effects on recreational activities.

Many states have imposed limitations on the amount of manure that may be applied to fields based on phosphorus soil test levels and expected crop removal of phosphorus. Figure 1 presents a general summary of phosphorus soil test levels that may limit crop production, soil test phosphorus ranges for manure allocation/disposal and phosphorus soil test values above which manure application is discouraged. At present, certain Kansas swine operations must operate under Kansas Department of Agriculture regulations pertaining to field phosphorus soil test values and swine manure application to agricultural lands.


## Manure Analysis

Average nutrient contents for various types of manure systems (table values) are often used to estimate nutrient credits for crop production, but have serious limitations in developing specific individual manure

**Figure 1. Phosphorus Management Model for Kansas Crop Production and Manure Management**



**Figure 2. Percent Of Inorganic Nitrogen Available To Crops For Various Manure Management Systems**



management plans. Determining plant-available nutrients from manure applications without analysis is difficult. Nutrient contents of specific animal manure systems are quite variable and are influenced by the type of animal involved, the ration fed, the manure storage and handling system (leaching and ammonia volatilization losses), and the moisture content of the manure. Additionally, allowances for differences in potential mineralization rates and volatilization losses cannot be reliably estimated without knowing the amounts of organic and inorganic nitrogen for a specific manure system.

Samples from individual manure systems should be submitted to a laboratory for analysis. At a minimum, manure samples should be analyzed for total nitrogen, organic nitrogen, inorganic ammonium nitrogen, total P<sub>2</sub>O<sub>5</sub> equivalent and total K<sub>2</sub>O equivalent determinations. Other nutrients and the salt/sodium hazard of the manure can also be requested but generally are not essential. For solid manure samples, most laboratories report pounds of nutrient per ton of manure on an as-received moisture basis. For liquid manure, results are generally reported as the pounds of nutrient per 1,000 gallons of manure or pounds of nutrient per acre-inch of lagoon water.

If only the total nitrogen in the manure is known (organic and inorganic nitrogen not determined by laboratory), Table 1 can be used to estimate the fraction of the total nitrogen in various manure systems expected to be present as organic and inorganic ammonium nitrogen. For example, if a producer is using solid dairy

manure containing 12 pounds total nitrogen / ton (organic and ammonium nitrogen content not determined), estimate that 45 percent of the total nitrogen is ammonium nitrogen (5.4 pounds per ton) and 55 percent of the total nitrogen would be organic nitrogen (6.6 pounds per ton). An estimate of the relative quantities of organic and inorganic nitrogen is needed to estimate the effects of application management on potential crop nutrient credits. Though not as desirable as having a complete manure analysis with organic and inorganic nitrogen components broken out, knowing the total nitrogen content of the manure when calculating nitrogen credits is far better than not having an analysis at all and relying on average table values.

Average nutrient contents for various types of manure can be used if an analysis is not available – but the accuracy of the nutrients credited will be severely limited. Table 2 presents some average manure nutrient credit values for various manure systems. Keep in mind, however, that these average values are based on many different systems – and consequently result in much larger margins of error than if the actual analysis of the manure is known.

### Nitrogen Credits

In general during the growing season after application, about 25 percent of the organic nitrogen in solid manure, 30 percent of the organic nitrogen in liquid manure and 20 percent of the organic nitrogen in compost will undergo mineralization in warm season cropping systems. Mineralization rates likely will be somewhat greater in southern climates and slightly lower in northern regions since soil microbial activity is dependant on soil temperature. Less organic nitrogen from manure/compost will become available to cool season crops (wheat for example), since soils are cooler and microbial activity slower during the primary period of crop nutrient uptake. About one-half or less of the amount of organic nitrogen mineralized during the first growing season after manure application will be mineralized during the second growing season. Even less nitrogen will be mineralized during the third growing season (Table 3).

**Table 1. Typical Ratio of Ammonium-Nitrogen and Organic-Nitrogen in Manure Systems<sup>1</sup>**

	Solid Manure Systems		Liquid Manure Systems	
	NH <sup>+</sup> <sub>4</sub> -N % of Total Manure N	Organic N	NH <sup>+</sup> <sub>4</sub> -N % of Total Manure N	Organic N
Dairy	45	55	50	50
Beef	35	65	50	50
Swine	60	40	70	30
Turkey	65	35	--	--

**Table 3. Estimated Percentage Of Organic Nitrogen Available To Crops After Manure Application**

	Year 1	Year 2	Year 3
	- - - - % N mineralized - - - -		
<b>Liquid Manure</b>	30	12	6
<b>Solid Manure</b>	25	12	6
<b>Compost</b>	20	6	3

All of the inorganic ammonium nitrogen in manure is potentially available to growing crops and is dependant on overall application management. Nearly all of the inorganic nitrogen in manure is present as ammonium-N. Very little nitrate-N is normally detected. The amount of inorganic ammonium-N ultimately available to the growing crop is greatly affected by several aspects of application management. Application method (broadcast, knife injected, sweep injected, etc.) and time of delay between broadcast application and incorporation have large effects on potential nitrogen loss of inorganic nitrogen from manure. Little or no nitrogen loss occurs with subsurface applications or immediately incorporated surface broadcast ma-

nure, while nearly all of the inorganic-N is lost for unincorporated surface manure applications.

Surface applications of manure are subject to severe losses of ammonia-N via volatilization under warm conditions. Even if manure applications are incorporated on the day of application, significant volatilization losses (10 to 20 percent) are likely to occur, while a delay

of only 24 hours will likely result in a loss of more than one-third the ammonium-N present (Figure 2). If incorporation is delayed one week or longer, essentially all of the ammonium nitrogen present may be lost through volatilization. Application of manure under cool conditions will result in a slower rate of volatilization loss, but the total loss will eventually be similar regardless of environmental conditions.

Injection of liquid manure below the soil surface is highly recommended as a way to minimize odor potential and increase nitrogen efficiency (eliminate volatilization losses). However, some research has indicated significant losses (10 to 30 percent) of inorganic nitrogen if knife

*Continued on page 8*

**Table 2. Representative Analysis Of Several Manure Systems – Animal Types<sup>1</sup>**

	Bedding Or Litter	Solid Manure Analysis (as received /wet sample analysis)				Total K <sub>2</sub> O
		% Dry Matter	Total N	NH <sub>4</sub> <sup>+</sup> -N <i>Lbs / ton</i>	Total P <sub>2</sub> O <sub>5</sub>	
Dairy	No	18	9	4	4	10
Dairy	Yes	21	9	5	4	10
Beef	No	15	11	4	7	10
Beef	Yes	50	21	8	18	26
Swine	No	18	10	6	9	8
Swine	Yes	18	8	5	7	7
Poultry	No	45	33	26	48	34
Poultry	Yes	75	56	36	45	34
Turkey	No	22	27	17	20	17
Turkey	Yes	29	20	13	16	13

	Bedding Or Litter	Liquid Manure Analysis (as received sample analysis)				Total K <sub>2</sub> O
		% Dry Matter	Total N	NH <sub>4</sub> <sup>+</sup> -N <i>Lbs / ton</i>	Total P <sub>2</sub> O <sub>5</sub>	
Swine	Pit	4	36	26	27	22
Swine	Lagoon	~1	4	3	2	4
Beef	Pit	11	40	24	27	34
Beef	Lagoon	~1	4	2	9	5
Dairy	Pit	8	24	12	18	29
Dairy	Lagoon	~1	4	2.5	4	5

<sup>1</sup> Table 1 and 2 values adapted from "Livestock Manure Management," M. Vitosh, H. Person and E. Purkhiser, Michigan State University Extension Bulletin, "Utilization of Animal Manure as Fertilizer," A. Sutton, D. Nelson, and D. Jones, Purdue University and "Vermont Manure Nutrient Management," B. Jokela, J. Rankin and S. Hawkins, University of Vermont, in "Computer Programs to Assist with Manure Nutrient Management," SSSA Symposium, Oct. 30-31, 1995.

## Example Solid Manure Nutrient Credit Worksheet

This worksheet is used to calculate the amount of crop available nutrients to credit warm season crops in the year of manure application. For solid manure, most laboratories report the amount of nutrients on an as-received moisture basis (pounds nutrient per ton), while the nutrient contents for liquid manure systems are normally reported on a thousand gallon or acre-inch basis (pounds nutrient per 1,000 gallons or acre-inch). Once the amount of manure nutrients available for the crop is estimated and the amount of nutrients required for the crop production system is determined then the amount of manure to uniformly apply can be calculated.

### Solid Manure Example

#### Solid Beef Manure Analysis

Total N – 10 lbs/ton

Organic N – 6 lbs/ton      Ammonium N – 4 lbs/ton

Total P<sub>2</sub>O<sub>5</sub> – 8 lbs/ton      Total K<sub>2</sub>O – 12 lbs/ton

#### Management Information

P Soil Test - Low

Broadcast Manure Application

Incorporated Day after Application

1. Estimate 65% of ammonium available to crop – 35% volatilization loss (from Figure 2).  
 $4 \text{ Lbs ammonium N / ton} \times 65\% = 2.6 \text{ lbs available ammonium N / ton}$
2. Credit 25% of organic N available in year of application (from solid manure worksheet)  
 $6 \text{ Lbs organic N / ton} \times 25\% = 1.5 \text{ lbs available organic N / ton}$
3. Total organic N credit and ammonium N credit for total N credit  
 $2.6 \text{ lbs ammonium N} + 1.5 \text{ lbs organic N} = 4.1 \text{ Lbs total available N / ton}$
4. Credit 50% of total P<sub>2</sub>O<sub>5</sub> available (very low, low, medium P soil test – from worksheet)  
 $8 \text{ lbs P}_2\text{O}_5 / \text{ton} \times 50\% = 4.0 \text{ lbs available P}_2\text{O}_5 / \text{Ton}$
5. Credit 85% of total K<sub>2</sub>O available (from worksheet)  
 $12 \text{ lbs K}_2\text{O} / \text{ton} \times 80\% = 10.2 \text{ lbs available K}_2\text{O} / \text{ton}$

**Table 4. Solid Manure Nutrient Crediting Worksheet**


**Example 1. (from preceding page)**

	Manure Lab Results	×	Nutrient Availability Factor	=	Plant Available Nutrients
	(lbs/ton)				(lbs/ton)
Organic N	6		25% Available In Year Of Application		1.5 Organic N
NH <sub>4</sub> <sup>+</sup> -N	4		NH <sub>4</sub> <sup>+</sup> -N Availability Factor From Fig. 2		2.6 NH <sub>4</sub> <sup>+</sup> -N
Total N	10				4.1 Sum Of NH <sub>4</sub> <sup>+</sup> -N & Organic N
Total P <sub>2</sub> O <sub>5</sub>	8		50% 50% for Very Low to Low P Soil Tests 100% for Medium to Very High P Soil Tests		4.0 Available P <sub>2</sub> O <sub>5</sub>
Total K <sub>2</sub> O	12		85% Potassium Efficiency Factor		10.2 Available K <sub>2</sub> O


**Your Farm**

	Manure Lab Results	×	Nutrient Availability Factor	=	Plant Available Nutrients
	(lbs/ ton)				(lbs/ton)
Organic N	_____		25% Available In Year Of Application		_____ Organic N
NH <sub>4</sub> <sup>+</sup> -N	_____		NH <sub>4</sub> <sup>+</sup> -N Availability Factor From Fig. 2		_____ NH <sub>4</sub> <sup>+</sup> -N
Total N	_____				_____ Sum Of NH <sub>4</sub> <sup>+</sup> -N & Organic N
Total P <sub>2</sub> O <sub>5</sub>	_____		50% for Very Low to Low P Soil Tests 100% for Medium to Very High P Soil Tests		_____ Available P <sub>2</sub> O <sub>5</sub>
Total K <sub>2</sub> O	_____		85% Potassium Efficiency Factor		_____ Available K <sub>2</sub> O

**Figure 1. Phosphorus Management Model for Kansas Crop Production and Manure Management**



**Figure 2. Percent Of Inorganic N Available To Crops For Various Manure Management Systems**



## Liquid Manure Nutrient Credit Worksheet

This worksheet is used to calculate the amount of crop available nutrients to credit warm season crops in the year of manure application. For liquid manure, most laboratories report the amount of nutrients on a thousand gallon or acre-inch basis (pounds nutrient per 1,000 gallons or acre-inch) while solid manure is normally reported on an as-received moisture basis (pounds nutrient per ton). Once the amount of manure nutrients available for the crop is estimated and the amount of nutrients required for the crop production system is determined then the amount of manure to uniformly apply can be calculated.

### Liquid Manure Example

#### Liquid Swine Manure Analysis

Total N – 4 lbs/1,000 gal

Organic N – 2 lbs/1,000 gal

Total P<sub>2</sub>O<sub>5</sub> – 3 lbs/1,000 gal

Ammonium N – 2 lbs/1,000 gal

Total K<sub>2</sub>O – 3.4 lbs/1,000 gal

#### Management Information

P Soil Test - High

Knife Injected Manure Application

Maintenance P Application Desired

1. Estimate 90% of ammonium available to crop – 10% loss (from Figure 2).  
 $2 \text{ lbs ammonium N} / 1,000 \text{ gal} \times 90\% = 1.8 \text{ lbs available NH}_4\text{-N} / 1,000 \text{ gal}$
2. Credit 30% of organic N available in year of application (from liquid manure worksheet)  
 $2 \text{ lbs organic N} / 1,000 \text{ gal} \times 30\% = 0.6 \text{ lbs available organic N} / 1,000 \text{ gal}$
3. Total organic N credit and ammonium N credit for total N credit  
 $1.8 \text{ lbs ammonium N} + 0.6 \text{ Lbs organic N} = 2.4 \text{ Lbs total available N} / 1,000 \text{ gal}$
4. Credit 100% of total P<sub>2</sub>O<sub>5</sub> available (high, very high soil P test – from worksheet)  
 $3 \text{ lbs P}_2\text{O}_5 / 1,000 \text{ gal} \times 100\% = 3.0 \text{ lbs available P}_2\text{O}_5 / 1000 \text{ gal}$
5. Credit 85% of total K<sub>2</sub>O available (from worksheet)  
 $3.4 \text{ lbs K}_2\text{O} / 1,000 \text{ gal} \times 85\% = 2.9 \text{ lbs available K}_2\text{O} / 1000 \text{ gal}$

**Table 5. Liquid Manure Nutrient Crediting Worksheet**


**Example 2. (from preceding page)**

	Manure Lab Results	×	Nutrient Availability Factor	=	Plant Available Nutrients
	(lbs/1,000 gallons)				(lbs/1,000 gallons)
Organic N	2.0		30%	Available In Year Of Application	0.6 Organic N
NH <sub>4</sub> <sup>+</sup> -N	2.0		90%	NH <sub>4</sub> <sup>+</sup> -N Availability Factor From Fig. 2	1.8 NH <sub>4</sub> <sup>+</sup> -N
Total N	4.0				2.4 Sum Of NH <sub>4</sub> <sup>+</sup> -N & Organic N
Total P <sub>2</sub> O <sub>5</sub>	3.0	100%	50% for Very Low to Low P Soil Tests 100% for Medium to Very High P Soil Tests		3.0 Available P <sub>2</sub> O <sub>5</sub>
Total K <sub>2</sub> O	3.4	85%	Potassium Efficiency Factor		2.9 Available K <sub>2</sub> O


**Your Farm**

	Manure Lab Results	×	Nutrient Availability Factor	=	Plant Available Nutrients
	(lbs/1,000 gallons)				(lbs/1,000 gallons)
Organic N	_____		30%	Available In Year Of Application	_____ Organic N
NH <sub>4</sub> <sup>+</sup> -N	_____		_____	NH <sub>4</sub> <sup>+</sup> -N Availability Factor From Fig. 2	_____ NH <sub>4</sub> <sup>+</sup> -N
Total N	_____				_____ Sum Of NH <sub>4</sub> <sup>+</sup> -N & Organic N
Total P <sub>2</sub> O <sub>5</sub>	_____	_____	50% for Very Low to Low P Soil Tests 100% for Medium to Very High P Soil Tests		_____ Available P <sub>2</sub> O <sub>5</sub>
Total K <sub>2</sub> O	_____	85%	Potassium Efficiency Factor		_____ Available K <sub>2</sub> O

**Figure 1. Phosphorus Management Model for Kansas Crop Production and Manure Management**



**Figure 2. Percent Of Inorganic Nitrogen Available To Crops For Various Manure Management Systems**



### Continued from page 3

injection is used. Denitrification is thought to be at least partially responsible since the injection zone becomes waterlogged and contains high levels of organic material – conditions favoring microbial denitrification. Sweep injectors dilute the liquid manure with more soil and minimize denitrification losses while eliminating the potential for nitrogen volatilization losses (Figure 2).

### **Phosphorus Credits**

Both organic and inorganic phosphorus forms are present in manure. Phosphorus is not subject to volatilization loss like inorganic manure nitrogen, but small amounts of organic and inorganic phosphorus can be moved by surface runoff from manure in storage and from unincorporated surface manure applications. For phosphorus nutrient crediting purposes, 50 percent of the total  $P_2O_5$  present in the manure should be credited for fields having phosphorus soil test values in the very low and low ranges (situations where crop responses to applied phosphorus applications are expected). For fields with soil test values in the high to very high ranges, 100 percent of the  $P_2O_5$  present in the manure should be credited (Bray P-1 or Mehlich III of > 20 ppm or Olsen phosphorus of > 12 to 13 ppm).

Starter fertilizer applications often result in crop response regardless of soil test values or complementary broadcast applications; however, manure applications are not a substitute for starter phosphorus or potassium fertilizer applications.

### **Potassium Credits**

Potassium is not present in organic forms in manure; all is present in an inorganic, plant-available form. Consequently, potassium availability from manure is not related to mineralization rates or soil microbial activity.

**Dale F. Leikam**  
Agronomy Specialist  
Nutrient Management

**Ray E. Lamond**  
Agronomy Specialist  
Soil Fertility

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.

Publications from Kansas State University are available on the World Wide Web at: <http://www.oznet.ksu.edu>

Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. In each case, credit Dale F. Leikam and Ray E. Lamond, *Estimating Manure Nutrient Availability*, Kansas State University, January 2003.

### **Kansas State University Agricultural Experiment Station and Cooperative Extension Service**

MF-2562

January 2003

It is the policy of Kansas State University Agricultural Experiment Station and Cooperative Extension Service that all persons shall have equal opportunity and access to its educational programs, services, activities, and materials without regard to race, color, religion, national origin, sex, age or disability. Kansas State University is an equal opportunity organization. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, Extension Districts, and United States Department of Agriculture Cooperating, Marc A. Johnson, Director.

The total  $K_2O$  present in manure should theoretically be potentially available for crop uptake in the year of application. However, research frequently supports crediting only about 80 to 90 percent of the total  $K_2O$  from manure in the year of application. This may be a result of the inherent difficulty of uniformly applying manure in many systems.

### **Summary**

Wise and proper use of manure can improve overall crop production efficiency and profitability for many fields in Kansas. At the same time, proper application minimizes potential for phosphorus movement to Kansas surface waters or leaching of nitrogen to groundwater. Since a significant portion of crop nutrients found in manure are present in an organic form and are unavailable for plant uptake until converted to inorganic nutrient forms by soil microbes, the amount of nutrients ultimately used by plants in the year of application is difficult to predict.

Soil temperature and moisture affect the activity of soil microbes and are nearly impossible to predict with a high degree of accuracy. However, development of manure nutrient management plans that take these and other factors into account can improve overall manure management, crop production efficiency, and profitability. Careful attention to the composition of manure from specific animal systems, method of manure application and timing of manure incorporation have potentially large effects on the amounts of nutrients available during the growing season.

The information presented in this publication was adapted from work conducted in Kansas and other states. Information from all these sources varies to some degree, but was interpreted to best represent conditions and situations representative of Kansas and Kansas producers.