

## Water Quality

The earliest irrigation in Kansas may have been from about 1650 to the early 1700s in a Taos Indian village in what is now Scott County State Park. The “modern” era of irrigation might be considered to date to the 1880s with the organization of irrigation ditch companies that began building diversion works and a canal system along the Arkansas River (Erbart, 1969). Rapid expansion of Kansas irrigation occurred following WWII for a variety of reasons including political/societal will, technology, and readily available energy (Figure 1). The 1945 water appropriation act, which provides the basis for Kansas water law today, was designed to encourage development of water resources. With improvements in irrigation well drilling and pumping equipment and the development of the Hugoton natural gas well field, irrigation acreage increased rapidly using groundwater from the Ogallala Aquifer.


The reported number of irrigated acres in Kansas has been relatively stable for several decades (Figure 1), but in actuality there has been a loss of irrigated acres in the west (Region 1) and an increase of irrigated acres in central (Region 2) and eastern (Region 3) Kansas, as shown in Figure 2 and Table 1. For the period of 1989 to 2014, the state had about a 1 percent decrease in irrigated acres, but more than 200,000 acres dropped out of irrigated production in western Kansas. Within Region 1, Groundwater Management District (GMD) 3 had the largest loss, amounting to 140,000 acres, while GMD 1 had a reduction of almost 100,000 acres.

**Table 1:** Irrigated acres and change comparisons for 1989 and 2014. (KDA DWR Irrigation Water Use Reports).

	1989	2014	Change in acres
Kansas	3,098,830	3,067,133	-31,697
Region 1 (Western KS)	2,329,975	2,124,410	-205,565
Region 2 (Central KS)	716,480	855,536	139,056
Region 3 (Eastern KS)	52,375	87,187	34,812


  

Irrigated Acreage Shifts within Region 1			
	1989	2014	Change in acres
GMD 1	291,574	194,846	-96,728
GMD 3	1,572,470	1,432,463	-140,007
GMD 4	359,016	386,798	27,782
Remainder of Region 1	106,915	110,303	3,388




**Figure 1:** Irrigated acreage trends for Kansas. Early estimates are based on various surveys. Since 1989, the irrigated acreage numbers are reported by irrigators on their annual water use reports submitted to the Kansas Department of Agriculture.

Irrigated acreage and water use changes in GMD1 are even more dramatic than indicated by the 1989 to 2104 data, which show a one-third reduction in irrigated acres. The peak authorized acres and authorized water use at the time of formation of GMD 1 in 1973 was around 425,000 acres and almost 700,000 acre-feet (ac-ft) of authorized water withdrawals (Figure 3) (Personal communication, GMD 1).



**Figure 2:** Irrigated acreage trend in Kansas by regions, 1989 to 2014.




**Figure 3:** Irrigated acreage trend in Western Kansas GMD 1, 1989 to 2014 and 1973 authorized acres and water withdrawals.

### System Type Acreage Trends

Irrigation system types have changed over time, switching from predominately surface flood irrigation to sprinkler irrigation, which is predominately center pivots (Figure 4). In 1970, about 18 percent of the 1.8 million irrigated acres were sprinkler irrigated. The volatility in the reported total irrigation acreage base until 1989 was due to the fact that irrigation data being reported in the annual water use report was based on authorized acres as opposed to acres actually irrigated. Nevertheless, much of the increase in the total irrigation acreage base during the 1970s was associated with the adoption of center pivot irrigation, with an increase of nearly an additional million acres irrigated by center pivots. In 1990, about one half of all acres were center pivot irrigated. Since 1990, the total irrigated acreage base has remained relatively stable, but center pivot irrigation now irrigates nearly 90 percent of all irrigated land.


Subsurface drip irrigation (SDI) irrigated acres are also shown in Figure 2 but represent less than 1 percent of the irrigation base. SDI was not reported as a separate system type until 2004, so early SDI acreage estimates were based on contractor surveys and appear to have been over-estimating SDI development. In 2003, the estimated SDI acreage, based on a producer survey, indicated approximately 14,000 acres. The recent data is based on the reported acres from the annual water use reports. “SDI combo” indicated that a field was irrigated by an SDI system and some other system type. An example could be SDI irrigated corners of a center pivot system.



**Figure 4:** Acreage trend of total irrigated acres, sprinkler system acres, and subsurface drip irrigation (SDI) acres in Kansas.

### Crop Acreage Trends

Irrigated corn production occurs on over one-half of all irrigated acres in Kansas, with the highest acreage peak occurring in 2013 at almost 2.0 million acres of the approximately 3.1 million irrigated acres for the state (Figure 5). Crop acreages are estimated since many irrigated fields are split between two or more crops. Other irrigated crops reported in 2014 include horticultural type crops such as golf/sports fields, truck farms, orchards, sod/turf farms, nursery, and grapes and other field crops, such as cotton, sunflower, oats, barley, rye, dry bean, and pasture.




**Figure 5:** Irrigated acreage of the top five irrigated crops in Kansas.

## Irrigation Water Diversion Trends


The total yearly amount of irrigation water diverted in Kansas is shown in Figure 6. A regression line through the reported pumping amount indicates a reduction in the total amount of water over the years, although the correlation is weak. The amount of precipitation that occurs during a given year also influences the annual diversion. In general terms, the 1990s were relatively wet years, while the 2000s were relatively dry. One of the highest rainfall years on record was 1993, while 2002 was one of the lowest, accounting for the corresponding valley and peak in water use, respectively. Note a secondary peak in water use during the nearly statewide drought years of 2011 and 2012. The conversion of flood irrigated land to center pivot irrigated land also continued during this time, increasing from roughly 50 percent to over 90 percent center pivot irrigated during those years. A center pivot system would generally have higher irrigation efficiency than a flood system.

There are other factors that could contribute to reduce pumping over time. The continuing decline of water table levels and a subsequent decrease in well yield could also contribute to reduced total water diversion. Tillage practices have also shifted over time as well. Reduced and no-till tillage systems are dominant in irrigated agriculture. Flood systems relied on clean furrows to distribute irrigation water across the field. Since most fields are now sprinkler irrigated, the need to reduce the amount of surface residue to establish good furrows is eliminated. Reduced tillage also reduces soil water losses due to soil disturbance and enhances precipitation capture and stored soil water retention. Higher residues also reduce early season soil evaporation losses. Adoption of improved irrigation management practices, such as irrigation scheduling, increased pumping depths, and subsequent increase in irrigation pumping costs could also play a role in reducing application depths over time.



**Figure 6:** Total irrigation water diverted and average application depth by year for Kansas.


The application depth by regions is shown in Figure 7. Region 1 roughly represents the western one-third of Kansas; Region 2, the middle third; and Region 3, the eastern third. Most of the irrigated acres are in Region 1, which also has the largest net crop irrigation requirements, so Region 1 and total for the state are very similar.



**Figure 7:** Regional average irrigation application depths by year for Kansas

## Crop Yield Trends

The four major seed crops grown in Kansas have had upward trends in yield as shown in Figures 8 through 11 for corn, soybean, grain sorghum, and wheat. The Kansas corn yield trend has had the most dramatic increase for both irrigated and dryland production, with irrigated corn yield improvements of approximately 2.1 bushels per acre for the period of record, which is over twice the dryland rate of 0.75 bushels per acre. The average irrigated yield increases for soybean and grain sorghum are 0.57 bu/ac and 0.54 bu/ac, respectively. Irrigated statewide estimates of soybean and grain sorghum ended in 2009. Irrigated wheat yield trend is 0.26 bu/ac. In 2009, the reported acres based on fallow or continuous wheat was switched to reporting of dryland (non-irrigated) wheat production.




**Figure 8:** Kansas Corn Yield Trends since 1974 (KDA Kansas Farm Facts).

## Irrigation Water Use Efficiency


Irrigation water use efficiency (IWUE) is the yield of a crop divided by the amount of irrigation water applied. Since yield has increased over time (Figures 8 to 11) and the average application depth has been trending downward, IWUE should be increasing. Southwest Kansas yield, irrigation application, and IWUE for corn are shown in Figure 12. IWUE has increased over the period of record by a value of 0.10 bushels per inch per year, although the correlation is weak due to high year-to-year variability.

**Southwest Kansas Corn Production, Irrigation Application and Irrigation Water Productivity**




**Figure 12:** Corn yield, irrigation application depth, and irrigation water use efficiency trends for southwest Kansas.

**Kansas Soybean Yield Trend**




**Figure 9:** Kansas Soybean Yield Trends, 1984 to 2009 (KDA Kansas Farm Facts).

**Kansas Grain Sorghum Yield Trend**



**Figure 10:** Kansas Grain Sorghum Yield Trends 1974 to 2009 (KDA Kansas Farm Facts).


**Kansas Wheat Yield Trend**



**Figure 11:** Kansas Wheat Yield Trends since 1974; reporting of format changed to irrigated and non-irrigated (dryland) practice in 2007. (KDA Kansas Farm Facts).

Seasonal variability is illustrated in Figure 13. This data was collected at the Southwest Kansas Research and Extension Center in Garden City, Kansas, and presented for illustrative purposes as it is well known that spatial variability of rainfall would exist across the region. The pan evaporation line is used as an indicator of crop water demand. More pan evaporation would correlate with sunny, warm, and windy days, which are the same conditions for high crop water use. Notice during the droughty 2000s years, the values were high. Annual rainfall and the in-season rainfall for July and August are also plotted. Annual rainfall during the 1990s was at or above normal as compared to several years below normal in the 2000s, especially 2002, which was about 6 inches below normal. Rainfall for the 2002 July/August period, however, was only slightly below normal. In addition, the in-season rainfall distribution is an important factor, the impact of which is not assessed with a simple seasonal or annual amount analysis. The drought years of 2011 and 2012 showed low in-season rainfall and below normal annual rainfall amounts. The pan evaporation values were the highest recorded for the period. Applied irrigation in 2011 was the highest since 1991, but conditions were

so extreme, irrigation corn yields were reduced as indicated by the low statewide yield value shown in Figure 8. Although yield can also be affected by heat, the loss of irrigation capacity over time likely also contributed to the loss of yield when combined with low precipitation and high crop water use need. The average application depth for the region for all crops and systems trended downward for the period of record, despite the wetter years being concentrated at the beginning of the period.



**Figure 13:** Irrigation, pan evaporation, and rainfall values for the SWREC, Garden City, Kansas.

### Irrigation System Efficiency

An abundance of literature is available on irrigation system efficiency values. There are many definitions of irrigation efficiency. Different terms are used to describe various aspects of irrigation, but much confusion, misuse, and misunderstanding of the concept is still possible. The Kansas State University Research and Extension Bulletin MF2243, “Efficiencies and Water Losses of Irrigation Systems” (available at <https://www.bookstore.ksre.ksu.edu/pubs/MF2243.pdf>) reviews the topic.

The general perception in Kansas is that surface irrigation systems have low irrigation efficiency, while center pivot systems with modern nozzle packages have high irrigation efficiency. While surface irrigation is not universally inefficient, the typical gated pipe furrow irrigation system in Kansas was probably operating at about 65 percent irrigation efficiency. That means about 65 percent of the water diverted to the field ended up in the managed root zone of the crop. The major loss for these systems was deep percolation (water drained below the managed root zone) and tailwater (water running off the end of the field). Furrow irrigation systems are labor intensive, so to minimize operation labor, long runs and fixed set times (12 or 24 hour intervals) were used, which may not have matched well to field characteristics. The

condition of the furrows, which change with irrigation and rainfall events, have a major influence on the advance of the water, and the irrigator would have to adjust accordingly. Correctly setting all gates to the proper discharge rate for the correct water advance rate down the furrow many times required as much art as science.

A four-year field demonstration project was conducted by the Northwest Kansas State University Research and Extension Center and the Northwest Kansas Groundwater Management District from 1978 through 1981. The results from this demonstration project are discussed to give some context to the changes that have occurred in irrigated agriculture and how irrigation water use has been affected. Field observations were made on a number of fields for the period, and records of average field yields and application depths were made. The data for the fields with corn production are shown in Figures 14 and 15.

At the time, flood and center pivot irrigation were roughly equal in acres, and few wells were equipped with water meters. The short-term goal of the project at the time was to observe and record current practices by installing water meters and soil water monitoring stations in each demonstration field. This was a demonstration project, so results between fields cannot be statistically compared. It is nevertheless interesting to note that the average yield across years for the two system types were essentially equal at 134.9 bushels per acre for the center pivots and 134.8 bushels for the surface flood systems, which may be somewhat indicated by the Figure 14 when examining the listing of the system type across the top of the chart. The average water application depth for the center pivots was 14.34 inches as compared to 21.02 inches for the flood systems. In Figure 15, the system type associated with the application depth tends to have the higher amounts associated with flood systems, although the highest application depth observed was applied by a pivot irrigator. While this is anecdotal evidence of a difference in irrigation efficiency between system types, the results are consistent with research literature summarized in the bulletin cited previously.

The shift from flood to center pivot sprinkler irrigation also allows irrigators to improve management efficiency. This stems from the ability to apply smaller increments of water during a given event, the irrigation frequency interval is decreased, and the irrigation event is interruptible. These features allow the irrigator to be able to better match application amounts to root zone soil water holding capacity and maintain the root zone soil water to more consistent soil water levels; in short — be better able to employ improved irrigation scheduling techniques.




for about one-third of the reduction in the average application depth. For fields that have low efficiency systems that result in a low irrigation capacity and deficit irrigation conditions, an increase in efficiency would allow the deficit to be reduced. Once deficit irrigated conditions are eliminated, then increased efficiency could result in reduced water application amounts.

Adoption of conservation tillage practices leaving high residue levels on the soil surface has the advantages of 1) not causing soil water loss due to the tillage operation, 2) improved capture of precipitation, and 3) reduced soil water evaporation. The application of irrigation water is also more easily accomplished on a center pivot irrigated field since the soil (furrow) is not used as part of the conveyance system for water across a field. Each of the factors can impact the water budget by increasing the available water supply to a crop. In addition, the center pivot system application depth and irrigation interval can be much more easily matched to the available root-zone soil water storage conditions.

For example, a flood irrigation system may need to apply 3 to 4 inches of water per event in order to achieve uniform water distribution across the field as compared

to a typical center pivot application depth of 1 inch. A flood system may take 10 to 14 days to complete an irrigation cycle as compared to 3 to 5 days for a center pivot. In order to prevent stress at the last watered part of the field, the irrigation event would need to start the number of days of the irrigation cycle in advance of the stress event at the end of the field. For the first irrigation, the root zone may still be small, so little soil water is available before stress begins. To prevent stress at the last irrigation zone, watering may have to begin in the first zone when little or no irrigation is needed. There may be less than one inch of root zone stage available when an irrigation of 3 or 4 inches must be started. With a reduced application amount and smaller irrigation cycle time, the likelihood of excess irrigation at the start of the irrigation cycle to prevent end of the cycle stress is minimal. A similar scenario occurs at the end of the season, when a full irrigation must be applied to satisfy a limited need to allow the crop to reach physiological maturity.

In conclusion, it would appear that the shift in irrigation system type from flood irrigation to center pivot irrigation has resulted in improved use of irrigation water in Kansas as indicated by the generally downward



**Figure 15:** Corn yield and irrigation application depth for various fields from the Northwest Kansas Water Conservation and Utilization Project arranged in order of irrigation application depth.

trend in application depth per acre. This is partially due to improved irrigation efficiency characteristics of the center pivot as compared to typical gated pipe furrow irrigation system in the state. This downward trend in application depths can also be partially associated with additional irrigation management and cultural practices that can be more easily adapted to sprinkler-irrigated fields. Since productivity in terms of yield appears to still be increasing, the improvements in irrigation and management efficiencies appear to still be offsetting productivity losses associated with loss of well capacities in many areas. Increase in corn productivity is also related

to improvements in corn hybrid development that was better adapted to local conditions.

**References:**

Erhart, A. 1969. Early Kansas Irrigation. One of three article series, "A page out of Irrigation History." Irrigation Age Magazine. Feb 1969. 4 pps.

USDA NASS. Various Years. Kansas Farm Facts. In cooperation with Kansas Department of Agriculture.

Kansas Irrigation Water Use Report. Various Years. Prepared in cooperation by the Kansas Water Office and the KDA, Division of Water Resources.

**Authors:**

Danny H. Rogers and Jonathan Aguilar

Extension Irrigation Engineers

K-State Research and Extension

Kansas State University, Department of Biological and Agricultural Engineering

Publications from Kansas State University are available at:  
<http://www.bookstore.ksre.ksu.edu>

Date shown is that of publication or last revision. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. In each case, credit Danny Rogers and Jonathan Aguilar, *Kansas Irrigation Trends*, Kansas State University, September 2017.

**Kansas State University Agricultural Experiment Station  
and Cooperative Extension Service**

K-State Research and Extension is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, Extension Districts, and United States Department of Agriculture Cooperating, John D. Floros, Director.

**MF2849 (Rev.) September 2017**