

# Hairy Vetch Toxicity

**Larry C. Hollis, DVM, M.Ag.**  
*Beef Veterinarian*

**Deon van der Merwe, BVS, Ph.D.**  
*Veterinary Toxicologist*



*Hairy vetch in midsummer.*  
Photo: The Ohio State University

Kansas producers have successfully used hairy vetch (*Vicia villosa*) as a forage for cattle grazing for years. But this palatable legume may cause major health problems, and even death, if cattle are not observed closely for signs of disease and removed immediately if they occur.

Signs initially include itching and scaly skin that is thickened to the touch, as well as diarrhea, conjunctivitis, and body wasting. The coat develops a rough appearance, with hair standing erect. Exudates may harden and form crusts on the skin surface. Cattle may appear to have photosensitization, but the entire body will be affected, not just the white or light-skinned areas. This disease is especially remarkable when it is observed in black-hided cattle.

Vetch-associated disease is most prevalent and severe in cattle 3 years old or older that have been grazing the plants for 10 to 20 days or longer. In some cases, disease problems do not appear for several weeks after grazing is initiated. Sickness rates can vary widely. Once signs become evident, recovery is not promising, with fatality rates usually running between 50 to 100 percent of affected animals. There is no known effective treatment.

It has been suggested that there may be a genetic susceptibility to vetch toxicity because Angus and Holsteins seem to be the breeds most commonly affected. Exactly what triggers the problem, or which toxic component of hairy vetch causes the disease, is unknown. The multi-systemic, granulomatous inflammatory changes observed microscopically by pathologists most closely resemble a type-IV hypersensitivity reaction. This suggests that cattle that have reacted and survived are far more likely to react even more severely if exposed to hairy vetch again.

## References

<http://www.kansasruralcenter.org/publications/hairyvetch.pdf>

<http://www.addl.purdue.edu/newsletters/2004/winter/vetch.asp>

Pancieria RJ, Mosier DA, Ritchey JW. Hairy vetch (*Vicia villosa* Roth) poisoning in cattle: update and experimental induction of disease. *J Vet Diagn Invest.* 1992 Jul;4(3):318-25.

Brand names appearing in this publication are for product identification purposes only.  
No endorsement is intended, nor is criticism implied of similar products not mentioned.

Publications from Kansas State University are available at [www.bookstore.ksre.ksu.edu](http://www.bookstore.ksre.ksu.edu)

Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. In each case, credit  
Larry C. Hollis, *Hairy Vetch Toxicity*, Kansas State University, October 2010.

**Kansas State University Agricultural Experiment Station and Cooperative Extension Service**

**MF-2948**

**October 2010**

---

K-State Research and Extension is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, Extension Districts, and United States Department of Agriculture Cooperating, Gary Pierzynski, Interim Dean and Director.